

CYTOLOGICAL DIAGNOSIS OF THYROID DISEASES: PRACTICAL SIGNIFICANCE AND MODERN APPROACHES

Ergasheva Gulshan Tokhirovna

Assistant of the Department of Clinical Sciences Asian International University, Bukhara,
Uzbekistan

E-mail: ergashevagulshantoxirovna@oxu.uz

<https://doi.org/10.5281/zenodo.15536117>

Abstract. *Cytological diagnosis of thyroid diseases plays a crucial role in modern endocrinology. Fine-needle aspiration biopsy (FNAB) offers high accuracy in verifying nodular formations, tumors, and autoimmune conditions, particularly in early stages. This article explores the methodology of specimen collection, cytomorphological criteria for evaluating cell samples, and the diagnostic value of this method in clinical decision-making.*

Keywords: *thyroid gland, cytology, FNAB, thyroid nodules, autoimmune thyroiditis, tumors.*

Introduction

The cytological diagnosis of thyroid diseases has become an indispensable part of modern endocrine pathology due to the increasing prevalence of thyroid nodules and autoimmune conditions in the global population. The thyroid gland, being highly sensitive to environmental, nutritional, and genetic factors, is prone to a wide range of disorders, including goiter, thyroiditis, benign adenomas, and malignant tumors. Many of these pathologies present with overlapping clinical symptoms, such as palpable nodules, hormonal imbalance, or nonspecific systemic complaints, making clinical assessment alone insufficient for an accurate diagnosis.

In this context, fine-needle aspiration biopsy (FNAB) has emerged as the gold standard method for preoperative morphological evaluation of thyroid lesions. It is a minimally invasive, cost-effective, and highly informative diagnostic procedure that allows the collection of cellular material for cytological examination. FNAB provides critical insights into the nature of thyroid nodules, differentiating between benign and malignant formations, and guiding the decision-making process regarding the necessity and extent of surgical intervention.

While imaging techniques such as ultrasonography (US), scintigraphy, and elastography play essential roles in detecting and characterizing thyroid lesions, they often fall short in determining the exact cytological or histological nature of the pathology. For instance, cold nodules on scintigraphy may or may not correspond to malignancy, and ultrasound features alone cannot distinguish between follicular adenoma and carcinoma due to their similar appearances. Therefore, microscopic evaluation of FNAB smears remains the definitive method for morphological verification, especially in nodular thyroid disease.

Furthermore, the integration of FNAB results with clinical, biochemical, and radiological data enhances diagnostic accuracy and allows for a more nuanced understanding of complex or coexisting pathologies, such as autoimmune thyroiditis combined with nodular hyperplasia or microcarcinoma. The importance of cytological diagnosis is especially evident in iodine-deficient regions, where nodular goiter is endemic, and the need for differential diagnosis is frequent.

Advancements in FNAB technique, including needle modifications and optimized smear preparation methods, have significantly increased the quality and informativeness of collected

material. In expert hands, the accuracy of FNAB in diagnosing thyroid malignancies can reach over 95%. However, challenges remain, particularly in cases of poorly cellular or bloody aspirates, and in interpreting borderline cytological categories.

Given the growing demand for precision medicine and personalized treatment planning, the role of thyroid cytology is expanding beyond initial diagnosis. It now contributes to monitoring disease progression, evaluating therapeutic response, and stratifying patients for surgical or conservative management. Consequently, cytologists, endocrinologists, and pathologists must develop an integrated approach to interpreting FNAB results within the broader clinical context.

This article aims to present a comprehensive overview of the methodology, morphological criteria, and diagnostic value of FNAB in thyroid diseases, highlighting its role in routine practice and its contribution to improving patient outcomes.

Methodology and FNAB Technique

Fine-needle aspiration biopsy of the thyroid is a minimally invasive method for extracting material using a fine needle. Modified needles with notches, as practiced at the Endocrinology Research Center (RAMS), improve sample collection and enable the acquisition of rich cellular material even without aspiration. Prepared smears are stained using the May-Grünwald–Giemsa technique and examined under a microscope.

The informativeness of the biopsy depends on the experience of the specialist and proper targeting of the puncture site. Poor cellular samples often do not yield a conclusive diagnosis, whereas rich smears allow for assessment of colloid composition, thyrocytes, proliferation signs, inflammation, or atypia.

Diagnostic Criteria for Goiter Disease (Expanded)

Goiter disease, or thyroid enlargement, represents a heterogeneous group of pathologies characterized by structural and functional changes in the thyroid gland. It may arise in response to iodine deficiency, autoimmune reactions, or other environmental and genetic influences. Cytologically, goiter is classified into three main morphological forms: diffuse, nodular, and mixed (diffuse-nodular). Each form demonstrates specific features in fine-needle aspiration biopsy (FNAB) smears that aid in establishing a reliable diagnosis and assessing disease activity.

1. Diffuse Goiter

Diffuse goiter is characterized by uniform enlargement of the thyroid gland without discrete nodular formations. Cytological smears typically reveal:

- Abundant dense colloid, often thick and viscous.
- Scattered flattened or cuboidal thyrocytes, usually without significant atypia.
- Low cellularity and absence of architectural complexity.

This type reflects colloid accumulation and gland hypertrophy in response to elevated thyrotropin (TSH) levels, often due to iodine deficiency.

2. Nodular Goiter

Nodular goiter (also referred to as multinodular goiter or nodular hyperplasia) involves the formation of one or more nodules within the thyroid tissue. These nodules may demonstrate varying degrees of:

- Follicular cell proliferation, sometimes forming microfollicular or papillary-like structures.
- Increased cellularity with small to moderately enlarged nuclei, occasional binucleated cells, and mild anisocytosis.

- The presence of regressive changes, such as:
 - Hemorrhage (with hemosiderin-laden macrophages or altered erythrocytes).
 - Cystic degeneration with proteinaceous or colloid fluid.
 - Fibrosis and hyalinization of nodule stroma.

Cytological evaluation must differentiate between benign nodular hyperplasia and neoplastic transformation, especially when cellular atypia or architectural complexity is present.

3. Mixed (Diffuse-Nodular) Goiter

This form combines features of both diffuse and nodular pathology. FNAB specimens may demonstrate:

- Heterogeneous background with colloid of varying consistency (dense, thin, or finely granular).
 - A mixture of resting follicular cells and active proliferative zones, showing microfollicular formations.
 - Possible coexistence of inflammatory infiltrates, particularly in cases with superimposed autoimmune components.
- This variant is commonly seen in patients with long-standing iodine deficiency or in those with evolving autoimmune thyroid disease superimposed on a multinodular background.

4. Cytological Indicators of Proliferative Activity

In both nodular and mixed forms of goiter, the following features point toward active follicular proliferation:

- Increased number of follicular epithelial cells, often forming clusters, microfollicles, or papillary-like projections.
- Cells with enlarged, hyperchromatic nuclei, but lacking definitive features of malignancy.
- Absence of mitotic figures, which distinguishes hyperplastic from neoplastic processes.
- Occasional nuclear crowding or overlapping, especially in hyperfunctioning nodules.

5. Importance of Colloid-Epithelial Balance

One of the key cytological criteria in diagnosing goiter is the ratio between colloid and epithelial elements:

- Predominantly colloid with sparse epithelial cells indicates colloid goiter.
- Rich cellularity with minimal colloid suggests proliferative (hyperplastic) goiter.
- Intermediate findings may reflect a transition or mixed state, requiring clinical correlation and, occasionally, follow-up FNAB.

6. Clinical and Ultrasound Correlation

Since cytological features alone may not always reveal the full spectrum of changes, especially in early or regressive stages, correlation with:

- Ultrasound findings (nodule size, echogenicity, vascularity),
 - Hormonal assays (TSH, T3, T4 levels), and
 - Clinical signs (growth pattern, compressive symptoms)
- is essential for comprehensive diagnosis and management.

In conclusion, accurate cytological assessment of goiter depends not only on identifying key cellular and colloidal features but also on understanding their biological context. Recognizing the morphological spectrum of goiter enables differentiation from follicular

neoplasms and assists in guiding appropriate clinical follow-up or intervention.

Cytological Features of Autoimmune Processes

Autoimmune thyroiditis (AIT) is characterized by lymphocytes, plasma cells, macrophages, and Askanazy (Hürthle) cells. Variants include classic (Hashimoto), lymphomatous, and fibrotic forms. Graves' disease shows abundant proliferating thyrocytes, cytoplasmic vacuolization, hypertrophic nuclei, and absence of colloid. The presence of peripheral blood in punctate reflects the gland's vascularity.

Benign and Malignant Neoplasms

Adenomas from A- and B-cells show varied structure (follicular, solid, papillary). Cytologically, they display high cellularity, clear cell boundaries, basophilic cytoplasm, and nuclear uniformity. Differentiation from nodular goiter is based on monomorphic proliferation.

Papillary carcinoma presents with nuclear grooves, pseudo-inclusions, and papillary structures. Follicular carcinoma resembles adenoma; definitive diagnosis is histological. Medullary carcinoma features granular cytoplasm, amyloid deposits, and metachromatic staining.

The Bethesda Classification System

The modern Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) includes six categories ranging from "non-diagnostic" to "malignant." This standardized approach improves communication between cytologists and clinicians and facilitates appropriate management decisions.

Practical Importance of Cytology

Cytological assessment of thyroid punctates allows for:

- avoiding unnecessary surgeries in benign cases;
- identifying malignancies that require surgery;
- monitoring nodular dynamics and treatment efficacy in AIT or Graves' disease;
- correlating findings with ultrasound and lab data for comprehensive evaluation

Conclusion

Cytological diagnosis of thyroid diseases is an essential tool in modern endocrinology. Its success depends on quality sampling, precise FNAB technique, and expert microscopic interpretation. Integrating cytology with clinical and imaging data enhances diagnostic accuracy and optimizes patient management outcomes.

REFERENCES

1. Брынова, О. В., Касоян, К. Т., Шабалова, И. П., Зима, А. П., Исаева, А. В., & Саприна, Т. В. (2016). Метод жидкостной цитологии в диагностике заболеваний щитовидной железы. *Клиническая лабораторная диагностика*, 61(4), 225-228.
2. Абенова, М. Т., Пикалова, Н. К., Жагапаров, М. К., Жумаканова, Ж. Т., & Токушпекова, Л. Б. (2011). Цитологическая диагностика заболеваний щитовидной железы. *Journal of Clinical Medicine of Kazakhstan*, (2 (21)), 144-146.
3. Ergasheva, G. (2025). POLYCYSTIC OVARY SYNDROME: A COMPREHENSIVE OVERVIEW AND CURRENT TREATMENT APPROACHES. *Modern Science and Research*, 4(4), 937-944.
4. Narzulaeva, U. R., & Bekkulova, M. A. (2023). Arterial gipertenziya etiologiyasida dislipidemiyaning xavf omili sifatidagi roli. *Science and Education*, 4(2), 415-419.

5. Ergasheva, G. (2025). ACROMEGALY: A SEVERE NEUROENDOCRINE DISORDER WITH MULTISYSTEM MANIFESTATIONS. *Modern Science and Research*, 4(3), 1123-1131.
6. Ergasheva, G. (2024). THE ROLE OF CORRECTIONAL PEDAGOGY IN ORGANIZING THE EDUCATION OF CHILDREN WITH DISABILITIES. *Ethiopian International Journal of Multidisciplinary Research*, 11(06), 206-207.
7. Zikrillaev, F. A. (2024). Cardiorehabilitations from Physiotherapeutic Treatments in Cardiovascular Diseases. *American Journal of Bioscience and Clinical Integrity*, 1(10), 96-102.
8. Зикриллаев, Ф. А. (2024). ОПРЕДЕЛЕНИЕ РАННИХ ФАКТОРОВ РИСКА ХРОНИЧЕСКОЙ БОЛЕЗНИ ПОЧЕК В ПУБЕРТНОМ ВОЗРАСТЕ. *Modern education and development*, 16(7), 166-180.
9. Toxirovna, E. G. (2024). QALQONSIMON BEZ KASALLIKLARIDAN HASHIMOTO TIREODIT KASALLIGINING MORFOFUNKSIONAL O'ZIGA XOSLIGI. *Modern education and development*, 16(7), 120-135.
10. Toxirovna, E. G. (2024). REVMATOID ARTRIT: BO'G'IMLAR YALLIG'LANISHINING SABABLARI, KLINIK BELGILARI, OQIBATLARI VA ZAMONAVIY DAVOLASH YONDASHUVLARI. *Modern education and development*, 16(7), 136-148.
11. Эргашева, Г. Т. (2024). ОЦЕНКА КЛИНИЧЕСКОЙ ЭФФЕКТИВНОСТИ ОРЛИСТАТА У БОЛЬНЫХ ОЖИРЕНИЕМ И АРТЕРИАЛЬНОЙ ГИПЕРТЕНЗИЕЙ. *Modern education and development*, 16(7), 92-105.
12. Ergasheva, G. T. (2024). THE SPECIFICITY OF AUTOIMMUNE THYROIDITIS IN PREGNANCY. *European Journal of Modern Medicine and Practice*, 4(11), 448-453.
13. Эргашева, Г. Т. (2024). ИССЛЕДОВАНИЕ ФУНКЦИИ ЩИТОВИДНОЙ ЖЕЛЕЗЫ ПРИ ТИРЕОИДИТЕ ХАШИМОТО. *Modern education and development*, 16(7), 106-119.
14. Toxirovna, E. G. (2024). GIPOFIZ ADENOMASINI NAZORAT QILISHDA KONSERVATIV JARROHLIK VA RADIATSIYA TERAPIYASINING UZOQ MUDDATLI SAMARADORLIGI. *Modern education and development*, 16(7), 79-91.
15. ERGASHEVA, G. T. (2024). OBESITY AND OVARIAN INSUFFICIENCY. *Valeology: International Journal of Medical Anthropology and Bioethics*, 2(09), 106-111.
16. Ergasheva, G. T. (2024). Modern Methods in the Diagnosis of Autoimmune Thyroiditis. *American Journal of Bioscience and Clinical Integrity*, 1(10), 43-50.
17. Tokhirovna, E. G. (2024). COEXISTENCE OF CARDIOVASCULAR DISEASES IN PATIENTS WITH TYPE 2 DIABETES. *TADQIQOTLAR. UZ*, 40(3), 55-62.
18. Toxirovna, E. G. (2024). DETERMINATION AND STUDY OF GLYCEMIA IN PATIENTS WITH TYPE 2 DIABETES MELLITUS WITH COMORBID DISEASES. *TADQIQOTLAR. UZ*, 40(3), 71-77.
19. Toxirovna, E. G. (2024). XOMILADORLIKDA QANDLI DIABET KELTIRIB CHIQUARUVCHI XAVF OMILLARINI ERTA ANIQLASH USULLARI. *TADQIQOTLAR. UZ*, 40(3), 63-70.

20. Toxirovna, E. G. (2024). QANDLI DIABET 2-TIP VA KOMORBID KASALLIKLARI BO'LGAN BEMORLARDA GLIKEMIK NAZORAT. *TADQIQOTLAR. UZ*, 40(3), 48-54.
21. Tokhirovna, E. G. (2024). MECHANISM OF ACTION OF METFORMIN (BIGUANIDE) IN TYPE 2 DIABETES. *JOURNAL OF HEALTHCARE AND LIFE-SCIENCE RESEARCH*, 3(5), 210-216.
22. Tokhirovna, E. G. (2024). THE ROLE OF METFORMIN (GLIFORMIN) IN THE TREATMENT OF PATIENTS WITH TYPE 2 DIABETES MELLITUS. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(4), 171-177.
23. Эргашева, Г. Т. (2024). Эффект Применения Бигуанида При Сахарным Диабетом 2 Типа И Covid-19. *Research Journal of Trauma and Disability Studies*, 3(3), 55-61.
24. Toxirovna, E. G. (2024). QANDLI DIABET 2 TUR VA YURAK QON TOMIR KASALLIKLARINING BEMOLARDA BIRGALIKDA KECISHI. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(7), 202-209.
25. Эргашева, Г. Т. (2024). СОСУЩЕСТВОВАНИЕ ДИАБЕТА 2 ТИПА И СЕРДЕЧНО-СОСУДИСТЫХ ЗАБОЛЕВАНИЙ У ПАЦИЕНТОВ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(7), 219-226.
26. Эргашева, Г. Т. (2024). СНИЖЕНИЕ РИСКА ОСЛОЖНЕНИЙ У БОЛЬНЫХ САХАРНЫМ ДИАБЕТОМ 2 ТИПА И СЕРДЕЧНО-СОСУДИСТЫМИ ЗАБОЛЕВАНИЯМИ. *Образование Наука И Инновационные Идеи В Мире*, 38(7), 210-218.
27. Tokhirovna, E. G. (2024). CLINICAL AND MORPHOLOGICAL ASPECTS OF THE COURSE OF ARTERIAL HYPERTENSION. *Лучшие интеллектуальные исследования*, 12(4), 234-243.
28. Tokhirovna, E. G. Studying the Causes of the Relationship between Type 2 Diabetes and Obesity. *Published in International Journal of Trend in Scientific Research and Development (ijtsrd)*, ISSN, 2456-6470.
29. Toxirovna, E. G. (2024). ARTERIAL GIPERTENZIYA KURSINING KLINIK VA MORFOLOGIK JIHATLARI. *Лучшие интеллектуальные исследования*, 12(4), 244-253.
30. Эргашева, Г. Т. (2024). НОВЫЕ АСПЕКТЫ ТЕЧЕНИЕ АРТЕРИАЛЬНОЙ ГИПЕРТОНИИ У ВЗРОСЛОГО НАСЕЛЕНИЕ. *Лучшие интеллектуальные исследования*, 12(4), 224-233.
31. Эргашева, Г. Т. (2024). ФАКТОРЫ РИСКА РАЗВИТИЯ САХАРНОГО ДИАБЕТА 2 ТИПА. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(5), 70-74.
32. Эргашева, Г. Т. (2024). ОСЛОЖНЕНИЯ САХАРНОГО ДИАБЕТА 2 ТИПА ХАРАКТЕРНЫ ДЛЯ КОГНИТИВНЫХ НАРУШЕНИЙ. *TADQIQOTLAR. UZ*, 30(3), 112-119.
33. Эргашева, Г. Т. (2023). Исследование Причин Связи Диабета 2 Типа И Ожирения. *Research Journal of Trauma and Disability Studies*, 2(12), 305-311.
34. Tokhirovna, E. G. (2024). Risk factors for developing type 2 diabetes mellitus. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(5), 64-69.

35. Toxirovna, E. G. (2024). QANDLI DIABET 2-TUR VA O'LIMNI KELTIRIB CHIQUARUVCHI SABABLAR. *Лучшие интеллектуальные исследования*, 14(4), 86-93.
36. Tokhirovna, E. G. (2023). Study of clinical characteristics of patients with type 2 diabetes mellitus in middle and old age. *Journal of Science in Medicine and Life*, 1(4), 16-19.
37. Toxirovna, E. G. (2024). GIPERPROLAKTINEMIYA KLINIK BELGILARI VA BEPUSHTLIKKA SABAB BO'LUVCHI OMILLAR. *Лучшие интеллектуальные исследования*, 14(4), 168-175.
38. Toxirovna, E. G. (2023). QANDLI DIABET 2-TUR VA SEMIZLIKNING O'ZARO BOG'LIQLIK SABABLARINI O'RGANISH. *Ta'lim innovatsiyasi va integratsiyasi*, 10(3), 168-173.
39. Saidova, L. B., & Ergashev, G. T. (2022). Improvement of rehabilitation and rehabilitation criteria for patients with type 2 diabetes.
40. Эргашева, Г. Т. (2023). Изучение Клинических Особенности Больных Сахарным Диабетом 2 Типа Среднего И Пожилого Возраста. *Central Asian Journal of Medical and Natural Science*, 4(6), 274-276.
41. Toxirovna, E. G. (2023). O'RTA VA KEKSA YOSHLI BEMORLARDA 2-TUR QANDLI DIABET KECHISHINING KLINIKO-MORFOLOGIK XUSUSIYATLARI. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 33(1), 164-166.
42. Ergasheva, G. T. (2022). QANDLI DIABET BILAN KASALLANGANLARDA REABILITATSIYA MEZONLARINI TAKOMILASHTIRISH. *TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMIY JURNALI*, 2(12), 335-337.
43. Ergasheva, G. (2024). METHODS TO PREVENT SIDE EFFECTS OF DIABETES MELLITUS IN SICK PATIENTS WITH TYPE 2 DIABETES. *Журнал академических исследований нового Узбекистана*, 1(2), 12-16.
44. ГТ, Э., & Саидова, Л. Б. (2022). СОВЕРШЕНСТВОВАНИЕ РЕАБИЛИТАЦИОННО-ВОССТАНОВИТЕЛЬНЫХ КРИТЕРИЕВ БОЛЬНЫХ С СД-2 ТИПА. *TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMIY JURNALI*, 2(12), 206-209.
45. Toxirovna, E. G. (2025). YURAK-QON TOMIR TIZIMI KASALLIKLARIDA BEMORLAR PARVARISHINING O'ZIGA XOSLIGI. *Modern education and development*, 20(2), 38-46.
46. Ergasheva, G. (2025). PECULIARITIES WHEN ACCOMPANIED BY HYPOTHYROIDISM AND IODINE DEFICIENCY IN PATIENTS WITH ADRENAL GLAND PATHOLOGY. *Modern Science and Research*, 4(2), 1133-1140.
47. Tokhirovna, E. G. (2024). Relationship of the Functional States of the Thyroid and the Reproductive System in Women under Iodine Deficiency. *Journal of Science in Medicine and Life*, 2(6), 89-94.
48. Зикриллаев, Ф. А. (2024). РОЛЬ ПРЕПАРАТОВ ЛЕРКАНИДИПИНА И АМЛОДИПИНА В ЛЕЧЕНИИ АРТЕРИАЛЬНОЙ ГИПЕРТЕНЗИИ ПРИ ХРОНИЧЕСКОЙ БОЛЕЗНИ ПОЧЕК. *Modern education and development*, 16(7), 213-229.

49. Abdurashitovich, Z. F. (2024). DETERMINATION OF THE ETIOPATHOGENESIS AND RISK FACTORS OF OBESITY AMONG ADOLESCENTS. *Modern education and development*, 16(7), 181-194.
50. Abdurashitovich, Z. F. (2024). EARLY RISK FACTORS FOR ARTERIAL HYPERTENSION AND FEATURES OF ITS ASSOCIATION WITH OTHER DISEASES. *Modern education and development*, 16(7), 195-212.