

THE RELATIONSHIP BETWEEN DIABETES MELLITUS AND THYROID DISORDERS: PATHOGENETIC MECHANISMS AND CLINICAL APPROACHES

Shaxloxon Rustamovna

Independent Researcher.

<https://doi.org/10.5281/zenodo.23281472>

Abstract. *Diabetes mellitus and thyroid disorders are common endocrine diseases that may frequently coexist and influence each other through complex metabolic, hormonal, and immunological mechanisms. The aim of this article is to analyze the relationship between diabetes mellitus and thyroid dysfunction, with particular attention to common pathogenetic mechanisms, clinical manifestations, diagnostic approaches, and principles of management.*

A review and analysis of scientific literature concerning the interaction between diabetes mellitus and thyroid disorders were performed. The findings indicate that thyroid dysfunction may significantly affect glucose metabolism, insulin sensitivity, lipid profile, and body weight in patients with diabetes mellitus. Hypothyroidism is mainly associated with reduced metabolic activity, weight gain, dyslipidemia, and increased insulin resistance, whereas hyperthyroidism may contribute to increased hepatic glucose production and impaired glycemic control.

The relationship is particularly important in patients with type 1 diabetes mellitus because of the common autoimmune mechanisms underlying pancreatic β -cell destruction and autoimmune thyroid diseases. Laboratory assessment, including thyroid-stimulating hormone and free thyroxine measurements, together with glycemic and metabolic parameters, may facilitate early identification of thyroid abnormalities.

Keywords: *Diabetes mellitus; thyroid dysfunction; hypothyroidism; hyperthyroidism; insulin resistance; thyroid hormones; glucose metabolism; autoimmune thyroid disease; type 1 diabetes mellitus; type 2 diabetes mellitus; endocrine disorders.*

Introduction

Diabetes mellitus and thyroid disorders are among the most common endocrine diseases worldwide and represent important public health problems due to their increasing prevalence and long-term complications. Although these conditions have traditionally been considered separate endocrine disorders, growing evidence indicates that they are closely interconnected through metabolic, hormonal, and immunological mechanisms. The coexistence of diabetes mellitus and thyroid dysfunction may complicate disease management, influence metabolic control, and increase the risk of cardiovascular and other systemic complications. Thyroid hormones, particularly thyroxine (T4) and triiodothyronine (T3), play an essential role in regulating energy expenditure, carbohydrate metabolism, lipid metabolism, protein synthesis, and tissue sensitivity to insulin. Consequently, abnormalities in thyroid function may affect glucose homeostasis and contribute to changes in insulin sensitivity. Hypothyroidism may be associated with weight gain, dyslipidemia, and reduced metabolic activity, while hyperthyroidism can increase glucose production and insulin requirements. These mechanisms demonstrate the important role of thyroid function in maintaining metabolic balance in patients with diabetes mellitus.

The relationship between diabetes and thyroid disorders is also influenced by autoimmune mechanisms. This association is particularly important in patients with type 1 diabetes mellitus, in whom autoimmune destruction of pancreatic β -cells may occur together with autoimmune thyroid diseases such as Hashimoto's thyroiditis and Graves' disease.

The presence of common genetic and immunological factors may increase the susceptibility to multiple autoimmune endocrine disorders. Therefore, early identification of thyroid dysfunction in patients with type 1 diabetes is considered an important component of comprehensive endocrine evaluation. In patients with type 2 diabetes mellitus, the relationship with thyroid dysfunction is more strongly associated with insulin resistance, obesity, chronic low-grade inflammation, and metabolic disturbances. Insulin resistance may influence thyroid function through alterations in hypothalamic-pituitary-thyroid regulation, while thyroid dysfunction can further contribute to disturbances in glucose and lipid metabolism. This bidirectional relationship may make glycemic control more difficult and potentially increase the risk of metabolic and cardiovascular complications.

Relevance

Diabetes mellitus and thyroid disorders are widespread endocrine diseases that frequently occur together and may have a significant impact on metabolic homeostasis and the overall clinical condition of patients. The interaction between these disorders is determined by complex hormonal, metabolic, and immunological mechanisms. Thyroid hormones influence carbohydrate and lipid metabolism, energy expenditure, and insulin sensitivity, while persistent disturbances in glucose metabolism may affect thyroid function and modify the clinical course of thyroid diseases.

Aim

The aim of this article is to analyze the relationship between diabetes mellitus and thyroid disorders, focusing on their common pathogenetic mechanisms, metabolic and immunological interactions, clinical manifestations, and modern approaches to diagnosis, monitoring, and treatment.

Main part

Diabetes mellitus and thyroid disorders share several interconnected pathogenic mechanisms that can influence the development and progression of both conditions. The relationship between these endocrine disorders is mediated by hormonal, metabolic, and immunological processes. Thyroid hormones, particularly triiodothyronine (T3) and thyroxine (T4), regulate basal metabolic activity and influence carbohydrate, lipid, and protein metabolism.

Changes in thyroid hormone concentrations can therefore affect glucose homeostasis and insulin sensitivity. Hypothyroidism is frequently associated with reduced metabolic activity, weight gain, dyslipidemia, and impaired insulin sensitivity. In contrast, excessive thyroid hormone activity may increase hepatic glucose production and peripheral insulin requirements.

Insulin resistance may also affect thyroid function through alterations in cellular metabolism and hypothalamic-pituitary-thyroid regulation. In addition, chronic hyperglycemia can contribute to metabolic disturbances that influence endocrine function. Autoimmune mechanisms are particularly important in patients with type 1 diabetes mellitus, where autoimmune destruction of pancreatic β -cells may coexist with autoimmune thyroid diseases.

Common genetic susceptibility and immune dysregulation may contribute to the simultaneous development of these conditions.

In type 2 diabetes, obesity, visceral adiposity, chronic low-grade inflammation, and insulin resistance may represent important links between diabetes and thyroid dysfunction.

Thyroid hormones play a fundamental role in maintaining metabolic homeostasis and regulating the utilization of carbohydrates, lipids, and proteins.

Triiodothyronine is the biologically more active thyroid hormone and has a major influence on cellular energy expenditure and metabolic activity. Thyroid hormones regulate hepatic glucose production, intestinal glucose absorption, and peripheral glucose utilization.

They also influence insulin secretion and the sensitivity of peripheral tissues to insulin. In hypothyroidism, reduced thyroid hormone activity may decrease energy expenditure and alter glucose utilization, potentially contributing to insulin resistance. Changes in lipid metabolism are also commonly observed, particularly an increase in total cholesterol and low-density lipoprotein cholesterol. These abnormalities may increase cardiovascular risk in patients who already have diabetes mellitus. Hyperthyroidism produces an opposite metabolic pattern characterized by increased energy expenditure, enhanced lipolysis, and increased hepatic glucose production.

Increased glucose turnover may result in deterioration of glycemic control in patients with diabetes. Thyroid hormones also affect adipose tissue function and mitochondrial activity, thereby influencing overall energy balance. The interaction between thyroid hormones and insulin signaling is therefore important for maintaining normal metabolic function. Disturbances in thyroid hormone levels can modify the clinical course of diabetes by changing insulin requirements and metabolic control.

The coexistence of diabetes mellitus and thyroid dysfunction can produce a complex clinical picture because the symptoms of these diseases may overlap. Patients with hypothyroidism may experience fatigue, weakness, weight gain, cold intolerance, dry skin, constipation, and reduced physical activity. Several of these manifestations can also occur in patients with poorly controlled diabetes or chronic diabetic complications. Hyperthyroidism, on the other hand, may present with weight loss, increased appetite, palpitations, sweating, anxiety, tremor, and heat intolerance. Increased thyroid hormone activity may accelerate glucose metabolism and complicate glycemic control in diabetic patients. In patients with type 1 diabetes mellitus, autoimmune thyroid disorders are of particular clinical importance because of their shared autoimmune background. Hashimoto's thyroiditis and Graves' disease may occur alongside type 1 diabetes and should therefore be considered during long-term follow-up. In type 2 diabetes mellitus, thyroid dysfunction may be associated with obesity, insulin resistance, dyslipidemia, and cardiovascular risk factors. The presence of thyroid dysfunction can modify metabolic parameters and make diabetes management more challenging. Conversely, persistent hyperglycemia may be associated with changes in thyroid function tests and may influence the interpretation of laboratory results. Clinical assessment should therefore include a detailed history, physical examination, and appropriate laboratory evaluation.

Accurate diagnosis of thyroid dysfunction in patients with diabetes mellitus requires a systematic evaluation of clinical symptoms and laboratory parameters. Thyroid-stimulating hormone is generally considered the primary laboratory marker for the initial assessment of thyroid function. Measurement of free thyroxine provides additional information and helps distinguish different forms of thyroid dysfunction. Free triiodothyronine may be useful in selected patients, particularly when hyperthyroidism is suspected. In patients with type 1 diabetes mellitus, assessment of thyroid autoantibodies can provide valuable information regarding the presence of autoimmune thyroid disease. Anti-thyroid peroxidase antibodies and anti-thyroglobulin antibodies are commonly evaluated when autoimmune thyroiditis is suspected. Glycemic status should simultaneously be assessed using fasting plasma glucose and glycated hemoglobin (HbA1c). Lipid profile evaluation may also be useful because thyroid dysfunction can influence cholesterol and triglyceride levels.

In selected cases, thyroid ultrasonography may provide additional structural information and help identify autoimmune or nodular changes. Screening strategies should be based on the patient's age, type of diabetes, clinical symptoms, family history, and risk factors for thyroid disease. Regular monitoring may be particularly relevant for patients with type 1 diabetes because of the increased association with autoimmune thyroid disorders. Early detection of subclinical thyroid dysfunction may allow timely intervention and reduce the risk of metabolic deterioration.

Management of patients with simultaneous diabetes mellitus and thyroid dysfunction requires an individualized and multidisciplinary approach. Treatment should address both glycemic control and the specific thyroid disorder while considering the patient's metabolic and cardiovascular risk profile. In patients with hypothyroidism, thyroid hormone replacement therapy is generally used to restore normal thyroid hormone levels and improve metabolic function. In patients with hyperthyroidism, treatment is selected according to the underlying cause and may include antithyroid medication, radioactive iodine therapy, or surgical intervention when clinically indicated. Diabetes management should be adjusted according to changes in metabolic status and insulin sensitivity. Correction of thyroid dysfunction may influence glucose requirements, making regular monitoring of blood glucose and HbA1c particularly important during treatment. Lifestyle modification also remains an important component of comprehensive management. A balanced diet, regular physical activity, appropriate body-weight management, and smoking cessation can contribute to improved metabolic health. In patients with obesity and insulin resistance, weight management may provide benefits for both diabetes and thyroid-related metabolic abnormalities.

Long-term monitoring should include assessment of thyroid function, glycemic control, lipid parameters, and cardiovascular risk factors. Patients with autoimmune endocrine disorders may require periodic evaluation because additional autoimmune conditions can develop over time. Effective communication between endocrinologists, primary care physicians, and other healthcare professionals can improve continuity of care.

Results

The analysis demonstrated a clinically significant relationship between diabetes mellitus and thyroid dysfunction, with metabolic, hormonal, and immunological mechanisms contributing to their coexistence. The reviewed evidence indicates that abnormalities of thyroid function may influence glucose metabolism and insulin sensitivity, while persistent disturbances in glucose homeostasis may also be associated with changes in thyroid function. Patients with hypothyroidism generally demonstrated metabolic characteristics including reduced energy expenditure, increased body weight, dyslipidemia, and impaired insulin sensitivity. These changes may contribute to difficulties in achieving optimal glycemic control, particularly in patients with established type 2 diabetes mellitus. The presence of hypothyroidism was also associated with an unfavorable metabolic profile, which may increase the overall cardiovascular risk of diabetic patients.

Hyperthyroid states showed a different metabolic pattern. Increased thyroid hormone activity was associated with enhanced energy expenditure, increased hepatic glucose production, and alterations in insulin requirements. In diabetic patients, these changes may contribute to fluctuations in blood glucose levels and make glycemic management more challenging. The findings therefore support the importance of considering thyroid status when unexplained changes in glycemic control occur.

A particularly important association was observed between type 1 diabetes mellitus and autoimmune thyroid disorders. The coexistence of these conditions can be explained, at least in part, by shared autoimmune mechanisms and genetic susceptibility.

Autoimmune thyroiditis and other thyroid abnormalities may develop during the course of type 1 diabetes, emphasizing the importance of periodic thyroid function assessment in these patients.

The analysis also demonstrated that thyroid dysfunction may be accompanied by changes in lipid metabolism. Patients with reduced thyroid function may develop elevated total cholesterol and low-density lipoprotein cholesterol levels. When combined with diabetes mellitus, these metabolic abnormalities may contribute to a greater cardiovascular risk and require more comprehensive monitoring. Laboratory assessment remains central to the identification of thyroid dysfunction in diabetic patients. Thyroid-stimulating hormone and free thyroxine measurements provide important information about thyroid functional status, while thyroid autoantibodies may assist in identifying autoimmune disease. Glycated hemoglobin, fasting plasma glucose, and lipid parameters provide complementary information about metabolic control.

Discussion

The findings of this analysis demonstrate a complex pathogenetic and clinical relationship between diabetes mellitus and thyroid disorders. The coexistence of these two groups of endocrine diseases may be associated with disturbances in metabolic homeostasis, hormonal regulation, and, in some cases, autoimmune processes. Therefore, assessment of thyroid function in patients with diabetes should not be considered merely an additional diagnostic procedure, but rather an important component of comprehensive endocrine evaluation. Thyroid hormones play a central role in the regulation of energy expenditure and metabolic activity. Reduced thyroid hormone availability may lead to decreased metabolic activity, weight gain, and abnormalities in lipid metabolism. When these changes occur together with insulin resistance, particularly in patients with type 2 diabetes mellitus, they may contribute to deterioration of metabolic control.

Consequently, the coexistence of hypothyroidism and type 2 diabetes may create additional challenges in achieving optimal glycemic and metabolic management.

Hyperthyroidism is characterized by an opposite metabolic pattern. Increased thyroid hormone activity may enhance hepatic glucose production and alter glucose utilization in peripheral tissues. As a result, patients with diabetes and hyperthyroidism may experience greater fluctuations in blood glucose levels and may require closer monitoring of their glycemic status.

These observations indicate that changes in thyroid function should not be evaluated independently from the clinical course of diabetes. Another important aspect of this relationship is the role of autoimmune mechanisms. In particular, type 1 diabetes mellitus is characterized by immune-mediated destruction of pancreatic β -cells and may coexist with autoimmune thyroid diseases. The presence of common genetic susceptibility and immune dysregulation may contribute to the development of conditions such as Hashimoto's thyroiditis and Graves' disease.

Therefore, long-term clinical follow-up and appropriate thyroid assessment are important in patients with type 1 diabetes.

Insulin resistance also represents an important link between diabetes and thyroid dysfunction.

Insulin resistance affects not only glucose metabolism but may also influence adipose tissue activity, inflammatory pathways, and hormonal regulation. Obesity and visceral adiposity can further intensify these metabolic abnormalities. For this reason, evaluation of thyroid function may be particularly relevant in diabetic patients with excessive body weight, metabolic syndrome, or persistent insulin resistance. The interaction between thyroid dysfunction and lipid metabolism is another clinically important consideration. Hypothyroidism may be accompanied by an unfavorable lipid profile, including increased levels of atherogenic lipoproteins. Since diabetes mellitus is itself an important risk factor for cardiovascular disease, the coexistence of diabetes and hypothyroidism may further increase the patient's overall cardiovascular risk. This emphasizes the importance of evaluating not only glycemic parameters but also lipid metabolism and other cardiovascular risk factors.

One of the major challenges in clinical practice is the overlap between symptoms of thyroid disorders and manifestations of diabetes or its complications. Fatigue, changes in body weight, reduced physical capacity, and alterations in cardiovascular function may occur in both conditions.

Therefore, clinical symptoms alone may not be sufficient to establish the presence of thyroid dysfunction. A combination of clinical assessment and laboratory testing, including TSH, free T4, and appropriate metabolic parameters, provides a more reliable basis for diagnosis. The findings also suggest that the coexistence of diabetes mellitus and thyroid dysfunction requires an individualized therapeutic approach. Correction of thyroid dysfunction may improve metabolic status, but changes in thyroid hormone levels during treatment can also affect glucose requirements. Therefore, patients undergoing treatment for thyroid disease should be monitored carefully for changes in blood glucose and other metabolic indicators.

Conclusion

The analysis confirms that diabetes mellitus and thyroid disorders are closely interconnected endocrine conditions with common metabolic, hormonal, and immunological mechanisms. Their coexistence may significantly influence glucose metabolism, insulin sensitivity, lipid profile, body weight, and overall cardiovascular risk. Thyroid dysfunction can affect glycemic control in patients with diabetes, while persistent metabolic disturbances associated with diabetes may also influence thyroid function. Hypothyroidism is commonly associated with weight gain, dyslipidemia, and increased insulin resistance, whereas hyperthyroidism may contribute to increased glucose production and fluctuations in blood glucose levels. The association between type 1 diabetes mellitus and autoimmune thyroid disorders is of particular clinical importance because of their shared immunological background.

Therefore, patients with type 1 diabetes require appropriate and regular assessment of thyroid function, especially when clinical symptoms or metabolic changes suggest possible thyroid dysfunction.

References

1. American Diabetes Association Professional Practice Committee. Prevention or Delay of Diabetes and Associated Comorbidities: Standards of Care in Diabetes-2026 // Diabetes Care. - 2026. - Vol. 49, Suppl. 1. - P. S50–S60.
2. American Diabetes Association Professional Practice Committee. Children and Adolescents: Standards of Care in Diabetes-2026 // Diabetes Care. - 2026. - Vol. 49, Suppl. 1. - P. S297–S338.

3. American Diabetes Association Professional Practice Committee. Glycemic Goals, Hypoglycemia, and Hyperglycemic Crises: Standards of Care in Diabetes-2026 // Diabetes Care. - 2026. - Vol. 49, Suppl. 1.
4. Biondi B., Kahaly G.J., Robertson R.P. Thyroid Dysfunction and Diabetes Mellitus: Two Closely Associated Disorders // Endocrine Reviews. - 2019. - Vol. 40, No. 3. - P. 789–824.
5. Díez J.J., Iglesias P. An analysis of the relative risk of thyroid disease in type 2 diabetes // Diabetes Care. - 2012. - Vol. 35, No. 2. - P. 372–378.
6. Palma C.C.S., et al. Prevalence of thyroid dysfunction in patients with diabetes mellitus // Diabetology & Metabolic Syndrome. - 2013. - Vol. 5. - Art. 58.
7. Kadiyala R., Peter R., Okosieme O.E. Thyroid dysfunction in patients with diabetes: clinical implications and screening strategies // International Journal of Clinical Practice. - 2010. - Vol. 64, No. 8. - P. 1130–1139.
8. Wang C. The relationship between type 2 diabetes mellitus and related thyroid diseases // Journal of Diabetes Research. - 2013. - Vol. 2013. - Art. 390534.
9. Udovcic M., Pena R.H., Patham B., Tabatabai L., Kansara A. Hypothyroidism and the Heart // Methodist DeBakey Cardiovascular Journal. - 2017. - Vol. 13, No. 2. - P. 55–59.